

Original Article

Tumour necrosis factor- α (TNF- α) and miRNA expression in frontal and temporal neocortex in Alzheimer's disease and the effect of TNF- α on miRNA expression in vitro

Doris Culpan, Patrick G Kehoe, Seth Love

Dementia Research Group, School of Clinical Sciences, University of Bristol, John James Buildings, Frenchay Hospital, Frenchay, Bristol, BS16 1LE, United Kingdom.

Received January 11, 2011; accepted March 21, 2011; Epub March 25, 2011; published May 15, 2011

Abstract: Micro-RNAs (miRNAs) are short non-coding RNAs capable of regulating gene expression at the translational level. A number of studies have suggested that the expression of several miRNAs is changed in AD. The pro-inflammatory cytokine tumour necrosis factor- α (TNF- α) is increased in serum and CSF in AD. We measured the expression of *TNFA* and several AD candidate gene-associated miRNAs (*let7a/b*, *miR-128a/b*, *miR-27a/b*, *miR-155*) in frontal and temporal neocortex from AD and control brains. The expression of these miRNAs was also measured after incubating non-differentiated (NDC) and retinoic acid -differentiated (DC) SH-SY5Y neuroblastoma cells with TNF- α . *TNFA* expression was similar in AD and control brains but *miR-128a/b* levels were significantly reduced in the temporal cortex and *miR-128b* in the frontal cortex in AD. MiRNA levels did not correlate with *TNFA* expression in brain tissue but exposure of NDC and DC SH-SY5Y cells to TNF- α caused a variable dose-dependent response in the level of some of the miRNAs studied. Our brain tissue findings argue against a role for TNF- α in influencing the expression of these miRNAs in AD.

Keywords: Alzheimer's disease, Tumour necrosis factor- α , microRNA, SH-SY5Y cells

Introduction

Micro-RNAs (miRNAs) are highly conserved, short non-coding single-stranded RNAs involved in translational regulation of messenger RNA (mRNA). They recognise and bind generally but not exclusively to the 3'-untranslated (3'UTR) of specific mRNAs and in doing so generally cause gene silencing [1,2]. Some miRNAs are reported to be brain-specific or brain-enriched [3-5]. The number of miRNAs expressed in human brain is higher than in other organs, most likely reflecting the greater diversity of cell types and subtypes [6,7].

Recently, changes in miRNA expression were identified in Alzheimer's disease (AD) that were thought to contribute to its development [8-11]. One study found altered expression of several miRNAs in brain tissue and cerebrospinal fluid (CSF) of early- and late-stage AD patients compared with controls [12]. Other studies sug-

gested that certain miRNAs up-regulate APP and BACE therefore potentially contributing to the overproduction of A β in AD [13-15].

The role of the immune system in the clearance of A β from the brain has been the subject of a number of recent studies [16-18]. Multiple cytokines are produced in response to A β -mediated activation of microglia and elevated levels of these cytokines have been found in serum and CSF from AD patients [16,17,19]. TNF- α is considered to be one of the most important pro-inflammatory cytokines produced by activated microglia and of major importance in AD. Elevated levels of TNF- α , encoded for by *TNFA* (OMIM:191160), appear to correlate with disease progression [20,21] and with the onset of cognitive deficits [22,23].

We measured miRNAs *let-7a*, *let-7b*, *miR-128a*, *miR-128b*, *miR-27a*, *miR-27b*, *miR-155* as well

as TNFA in brain tissue from patients with AD and age-matched controls. These miRNAs were selected for their reported abundant expression in brain, involvement in inflammation and microglial activation [24-26], implicated in cell cycle regulation [27,28] and apoptosis [29]. In addition, other predicted targets of miR-27a/b and miR-128a/b as identified by the online software 'TargetScanHuman release: 5.1' include ACE, ECE1 and ECE2, which influence cerebral blood flow and are also capable of degrading A β . We also investigated if TNF- α influences the expression of any of these miRNAs in non-differentiated (NDC) and retinoic acid - differentiated (DC) SH-SY5Y neuroblastoma cells.

Materials and methods

Human brain tissue

This retrospective case-control study used brain tissue that had been donated to the Human Tissue Authority (HTA) licensed South West Dementia Brain Bank (SWDBB), Bristol. Sections of post mortem brain tissue were collected in RNAlater (Ambion, UK) and frozen at -80°C until RNA extraction. mRNA and miRNA from frontal and temporal neocortex (Brodmann areas 6 and 22) was isolated from 6 control (73–94 y, mean 87 y; 5 male, 1 female) and 12 AD (67–97 y, mean 81 y; 5 male, 7 female). The AD cases had 'definite AD' according to CERAD criteria [30]. The study had local Research Ethics Committee approval.

Cell culture and TNF- α exposure

SH-SY5Y human neuroblastoma cells were cultured in Dulbecco's modified Eagles medium (DMEM) (Sigma) (2 mM glutamine (Sigma), 1% essential amino acids (Sigma), 15% fetal bovine calf serum (FCS) (Autogene Bioclear). Cells were differentiated by addition of 10 μ M retinoic acid (RA) (Sigma-Aldrich, Dorset, UK) for 5 days and 50 ng/ml BDNF for 2 days in DMEM. DC and NDC were incubated in serum-free medium for 12-14 h before exposure to 1 ng/ml, 10 ng/ml or 50 ng/ml TNF- α (R&D systems, UK) for 15 min and 3 h. All cells were discarded after 15 passages as the sensitivity of SH-SY5Y cells to TNF- α was shown to decrease with increased passages [31]. All experiments were repeated 3 times, and on each occasion the mean values were determined for triplicate samples.

mRNA/miRNA extraction and real-time PCR (RT-PCR)

mRNA/miRNA was extracted using the mirVANA extraction kit (Ambion) according to the manufacturer's instructions. cDNA was produced from the mRNA and miRNA by use of the High Capacity c-DNA Archive Kit (Applied Biosystems). RT-PCR was performed using the ABI 7000 sequence detection system (Applied Biosystems) with Assay-on-Demand Gene Expression Products for TNFA, GAPDH and TaqMan Universal PCR Master Mix. RT-PCR of miRNAs was performed using TaqMan miRNA assays, detecting the mature miRNAs, (hsa-let-7a/7b, hsa-miR-155, has-miR-128a/128b, has-miR27a/27b) (Applied Biosystems) as described by the manufacturer. Expression of TNFA relative to GAPDH (calibrator gene) mRNA and of the individual miRNAs relative to RNUB6 (internal normalizing control) was calculated by the $2^{-\Delta\Delta Ct}$ method [32]. The results were expressed as the fold difference in gene expression between AD and controls.

Statistical analysis

Statistical tests were performed using GraphPad Prism v5 for windows. Mann-Whitney test was used for the comparison of expression levels across the subject groups. Differences with a $p < 0.05$ were considered significant. Since mRNA and miRNA levels were expressed as an exponential function (the fold-change relative to the appropriate control measurements) the values from repeat analyses were presented as the geometric means and interquartile ranges.

Results

Expression of TNF- α and miRNAs in AD and control cases brain tissue

Expression of TNFA, let-7a/b, miR-128a/b, miR-27a/b and miR-155 was detected in temporal and frontal neocortex from all brain samples studied although expression varied considerably between cases in all diagnostic groups. TNFA levels were particularly variable (**Table 1**) but did not differ significantly between the AD and controls in either temporal or frontal neocortex.

Expression of miR-128a/b was significantly decreased in the temporal neocortex in AD (miR-

Table 1. Expression of *TNFA* and *miRNAs* in AD and control brain tissue

AD	Frontal neocortex		Temporal neocortex	
	2 ^{-$\Delta\Delta$Ct}	p-value	2 ^{-$\Delta\Delta$Ct}	p-value
Control=1	Geometric mean (IQR)		Geometric mean (IQR)	
<i>TNFA</i> relative to <i>GAPDH</i>	0.72 (0.9)	NS	0.83 (1)	NS
<i>Let7a</i>	0.9 (0.59)	NS	0.9 (0.77)	NS
<i>Let 7b</i>	0.95 (0.45)	NS	0.54 (0.46)	NS
<i>miR-128a</i>	0.92 (0.27)	NS	0.68 (0.24)	0.006
<i>miR-128b</i>	0.69 (0.11)	0.006	0.77 (0.28)	0.044
<i>miR-155</i>	0.95 (0.38)	NS	0.96 (0.98)	NS
<i>miR-27a</i>	0.7 (0.75)	NS	0.80 (0.47)	NS
<i>miR-27b</i>	0.66 (0.59)	NS	0.62 (0.37)	NS

Table 2. miRNA expression in NDC SH-SY5Y cells after exposure to TNF- α

15mins	<i>miR-128a</i> 2 ^{-$\Delta\Delta$Ct} geometric mean	<i>miR-128b</i> 2 ^{-$\Delta\Delta$Ct} geometric mean	<i>miR-27a</i> 2 ^{-$\Delta\Delta$Ct} geometric mean	<i>miR-27b</i> 2 ^{-$\Delta\Delta$Ct} geometric mean	<i>let-7a</i> 2 ^{-$\Delta\Delta$Ct} geometric mean	<i>let-7b</i> 2 ^{-$\Delta\Delta$Ct} geometric mean	<i>miR-155</i> 2 ^{-$\Delta\Delta$Ct} geometric mean
1 ngTNF- α	1.32	1.4	1.04	1.36	0.8	0.72	0.97
10ngTNF- α	1.12	1.66	1.2	0.77	1.43	0.87	0.74
50ngTNF- α	0.75	1.11	0.91	0.61	2	1.51	1.21
3 h							
1 ngTNF- α	2.2	0.59	0.96	1.05	0.91	1.55	0.95
10ngTNF- α	2.09	1.38	0.98	1.28	0.92	1.89	1.56
50ngTNF- α	1.33	1.33	1.08	1.14	1.58	1.62	1.15

128a p=0.006, miR-128b p=0.044) and miR-128b (p=0.006) in the frontal cortex compared to controls. (Table 1). There was no association between the expression of TNFA and any of the miRNAs studied in the frontal or temporal cortex in either of the cohorts.

miRNA expression in DC and NDC SH-SY5Y cells after TNF- α exposure

The effect of TNF- α exposure on miRNA gene expression in vitro is summarised in Tables 2 and 3. In NDCs, let-7a/b expression increased and miR-128a and miR-27b decreased in a dose-dependent manner after 15 min exposure to TNF- α . After 3 h incubation only let-7a/b and miR-128a showed a response to TNF- α (Table

2).

The expression of miR-155 in control DCs was significantly lower than in control NDCs (2^{-DDCt} = 0.339) and after exposure to TNF- α reduced to the minimum level of detection (Table 3). MiR-128b expression was 2-fold higher (2^{- $\Delta\Delta$ Ct} = 2.21) in control DCs compared to control NDCs and miR-128b expression was consistently reduced in DCs after 3h TNF-a.

Discussion

Some studies have reported that miRNA expression profiles are altered in AD which may have an aetiological basis in AD. TNFA expression in brain was generally low in all individuals studied irrespective of their diagnostic groups and there

Table 3. miRNA expression in DC SH-SY5Y cells after exposure to TNF- α

15mins	<i>miR-128a</i> 2- $\Delta\Delta$ Ct	<i>miR-128b</i> 2- $\Delta\Delta$ Ct	<i>miR-27a</i> 2- $\Delta\Delta$ Ct	<i>miR-27b</i> 2- $\Delta\Delta$ Ct	<i>let-7a</i> 2- $\Delta\Delta$ Ct	<i>let-7b</i> 2- $\Delta\Delta$ Ct	<i>miR-155</i> 2- $\Delta\Delta$ Ct
	geometric mean	geometric mean	geometric mean	geometric mean	geometric mean	geometric mean	geometric mean
1 ngTNF- α	0.53	1.13	1.11	1.07	0.03	0.44	0.07
10ngTNF- α	1.3	0.88	1.52	1.5	0.42	0.4	0.03
50ngTNF- α	2.0	1.08	1.81	2.39	1.32	0.78	0.04
3 h							
1 ngTNF- α	1.18	0.37	1.19	1.19	0.76	1.02	0.16
10ngTNF- α	1.2	0.38	0.74	0.85	0.57	1.31	0.11
50ngTNF- α	1	0.54	1.05	1.15	1.45	1.35	0.32

was considerable variation between individuals. The generally low levels of TNFA expression in AD and control tissue may reflect a short half-life of TNF- α mRNA in brain tissue and the absence of factors which normally regulate TNFA mRNA stability [33,34]. There was no difference between mean TNFA expression levels in tissue from AD when compared to control subjects, confirming results of a previous study [35]. There was also no evidence of a correlation between TNFA and miRNA gene expression in brain tissue.

We observed a statistically significant decrease in miR-128a/b in temporal neocortex from AD compared to control brains and reduced miR-128b in the frontal cortex of AD brains. In contrast to these findings, miR-128 was previously shown to be up-regulated in the hippocampus in AD [36]. Diverse expression of miRNAs in distinct areas of the brain has been previously noted, including substantial variation in relative miRNA levels across different regions [9]. Recently miRNA deregulation in response to A β exposure in hippocampal neurons and in the hippocampus of A β 42-depositing APP23 mice at the onset of A β plaque formation was reported. It was suggested this may be an important factor contributing to the cascade of events leading to AD [37].

Several other reports have indicated the importance of miR-128 in neurogenesis and AD: it plays an important role in the differentiation of neuronal stem cells (NSCs) into neurons or astrocytes, regulates tau degradation and adenosine 2B receptor expression [38-40]. However, further investigation is needed to identify if the decreased levels observed for miR-128a/b in AD causes translational changes affecting genes for which interactions have been vali-

dated or other potential target genes. This is a challenging task since each miRNA can bind multiple targets and many miRNAs can bind the same target [4,9].

It has been suggested that miRNAs are normally relatively stable [41] but that their expression profile may change rapidly, as demonstrated previously in primary hippocampal neurons in response to A β treatment [37] and as we observed in this study in neuroblastoma cells in response to TNF- α .

We noticed that RA-induced differentiation of SH-SY5Y neuroblastoma cells alone reduced the expression of miR-155 and increased the expression of miR-128b. The increased expression of miR-128 in RA-differentiated neuroblastoma cells was reported previously to impair cell growth [42]. The expression of several other miRNAs was reported to be altered in cells exposed to RA [43]. Although miR-155 was not mentioned in that report, our findings indicate that it can be included in the list of miRNAs sensitive to RA-induced cell differentiation.

We found that miR-155 expression in differentiated neurons was severely reduced by TNF- α as has been reported previously in immune cells [44]. MiR-155 is a multifunctional miRNA, involved in numerous biological processes including haematopoiesis, inflammation and immunity. Deregulation of miR-155 has been associated with cancer, cardiovascular disease and viral infections [25]. It also affects the expression of type 1 angiotensin II receptor (AT1R) and PICALM (phosphatidylinositol binding clathrin assembly protein) [45], molecules thought to be involved in AD [46,47].

miRNAs are generally regarded as negative

regulators of mRNA translation and repression of miRNAs would be expected to enhance mRNA translation of target genes. However, under certain stress conditions miRNAs increase translation of their targets [48]. Previous work showed that members of the let-7 family of miRNAs bound to the 3'UTR of TNFA mRNA and by doing so stimulated translation in cells that were deprived of serum and therefore in a particular phase of cell cycle arrest [48].

It is possible that the initial up-regulation of these miRNAs targets genes involved in the TNF-R signaling pathways leading to neuronal cell survival or cell death. Over-expression of a miRNA complex including miR-27a sensitized HEK293T cells to TNF- α cytotoxicity and miR-27a negatively regulated the expression of FADD (Fas Associated protein with Death Domain), a protein involved in cell apoptosis [29]. It seems likely that the high serum and CSF levels of TNF- α observed in AD will influence miRNAs in this disease in vivo. However, further work is needed to establish the full range of miRNAs that are affected and the biological relevance of their targets to AD.

Acknowledgements

This study was supported by equipment support grants from the Alzheimer's Research Trust, salary support grants by BRACE (Bristol Research into Alzheimer's and Care of the Elderly), and a HEFCE funded Research Fellowship.

Address correspondence to: Dr. Doris Culpan, Dementia Research Group, School of Clinical Sciences, University of Bristol, John James Building, Frenchay Hospital, Frenchay, Bristol, BS16 1LE, United Kingdom. Tel: 0117 3403068; Fax: 0117 957 3955; E-mail: Doris.Culpan@bristol.ac.uk

References

- [1] Ambros V. The functions of animal microRNAs. *Nature* 2004; 431: 350-355.
- [2] Bartel DP and Chen C-Z. Micromanagers of gene expression: the potentially widespread influence of metazoan microRNAs. *Nat Rev Genet* 2004; 5: 396-400.
- [3] Lagos-Quintana M, Rauhut R, Yalcin A, Meyer J, Lendeckel W and Tuschl T. Identification of Tissue-Specific MicroRNAs from Mouse. *Current biology* : CB 2002; 12: 735-739.
- [4] Sempere L, Freemantle S, Pitha-Rowe I, Moss E, Dmitrovsky E and Ambros V. Expression profiling of mammalian microRNAs uncovers a subset of brain-expressed microRNAs with possible roles in murine and human neuronal differentiation. *Genome Biology* 2004; 5: R13.
- [5] NELSON PT, HATZIGEORGIOU AG and MOURELATOS Z. miRNP:mRNA association in polyribosomes in a human neuronal cell line. *RNA* 2004; 10: 387-394.
- [6] Berezikov E, Thuemmler F, van Laake LW, Kondova I, Bontrop R, Cuppen E and Plasterk RHA. Diversity of microRNAs in human and chimpanzee brain. *Nat Genet* 2006; 38: 1375-1377.
- [7] Beuvink I, Kolb FA, Budach W, Garnier A, Lange J, Natt F, Dengler U, Hall J, Filipowicz W and Weiler J. A novel microarray approach reveals new tissue-specific signatures of known and predicted mammalian microRNAs. *Nucleic Acids Research* 2007; 35: e52.
- [8] Nelson PT and Keller JN. RNA in Brain Disease: No Longer Just "The Messenger in the Middle". *Journal of Neuropathology & Experimental Neurology* 2007; 66: 461-468. DOI: 10.1097/1001.jnen.0000240474.0000227791.f0000240473.
- [9] Nelson PT, Wang WX and Rajeev BW. MicroRNAs (miRNAs) in Neurodegenerative Diseases. *Brain Pathology* 2008; 18: 130-138.
- [10] Barbato C, Arisi, I, Frizzo, M E, Brandi, R, Da Sacco, L, Masotti, A. Computational Challenges in miRNA Target Predictions: To Be or Not to Be a True Target? *J Biomed Biotechnol* 2009; 2009: 803069.
- [11] Kocerha J, Faghihi MA, Lopez-Toledano MA, Huang J, Ramsey AJ, Caron MG, Sales N, Willoughby D, Elmen J, Hansen HF, Orum H, Kauppinen S, Kenny PJ and Wahlestedt C. MicroRNA-219 modulates NMDA receptor-mediated neurobehavioral dysfunction. *Proceedings of the National Academy of Sciences* 2009; 106: 3507-3512.
- [12] Cogswell John P WJ, Taylor Ian A, Waters Michelle, Shi Yunling, Cannon Brian, Kelnar Kevin, Kemppainen Jon, Brown David, Chen Caifu, Prinjha Rab K, Richardson Jill C., Saunders Ann M., Roses Allen D., Richards Cynthia A. Identification of miRNA changes in Alzheimer's disease brain and CSF yields putative biomarkers and insights into disease pathways. *JOURNAL OF ALZHEIMER'S DISEASE* 2008; 14: 27-41.
- [13] Hebert SS and De Strooper B. MOLECULAR BIOLOGY: miRNAs in Neurodegeneration. *Science* 2007; 317: 1179-1180.
- [14] Wang W-X, Rajeev BW, Stromberg AJ, Ren N, Tang G, Huang Q, Rigoutsos I and Nelson PT. The Expression of MicroRNA miR-107 Decreases Early in Alzheimer's Disease and May Accelerate Disease Progression through Regulation of β -Site Amyloid Precursor Protein-Cleaving Enzyme 1. *J. Neurosci.* 2008; 28: 1213-1223.
- [15] Boissonneault V, Plante I, Rivest S and Provost P. MicroRNA-298 and MicroRNA-328 Regulate

- Expression of Mouse β -Amyloid Precursor Protein-converting Enzyme 1. *Journal of Biological Chemistry* 2009; 284: 1971-1981.
- [16] Heneka MT and O'Banion MK. Inflammatory processes in Alzheimer's disease. *Journal of Neuroimmunology* 2007; 184: 69-91.
- [17] Boche D and Nicoll JA. SYMPOSIUM: Clearance of A β from the Brain in Alzheimer' Disease: The Role of the Immune System in Clearance of A β from the Brain. *Brain Pathology* 2008; 18: 267-278.
- [18] Sastre M, Walter J and Gentleman S. Interactions between APP secretases and inflammatory mediators. *Journal of Neuroinflammation* 2008; 5: 25.
- [19] McGeer P and McGeer E. Local neuroinflammation and the progression of Alzheimer's disease. *J Neurovirol* 2002; 8: 529 - 538.
- [20] Paganelli R, Di Iorio A, Patricelli L, Ripani F, Sparvieri E, Faricelli R, Iarlori C, Porreca E, Di Gioacchino M and Abate G. Proinflammatory cytokines in sera of elderly patients with dementia: levels in vascular injury are higher than those of mild-moderate Alzheimer's disease patients. *Experimental Gerontology* 2002; 37: 257-263.
- [21] Holmes C, Cunningham C, Zotova E, Woolford J, Dean C, Kerr S, Culliford D and Perry VH. Systemic inflammation and disease progression in Alzheimer disease. *Neurology* 2009; 73: 768-774.
- [22] Janelsins M, Mastrangelo M, Oddo S, LaFerla F, Federoff H and Bowers W. Early correlation of microglial activation with enhanced tumor necrosis factor-alpha and monocyte chemoattractant protein-1 expression specifically within the entorhinal cortex of triple transgenic Alzheimer's disease mice. *J Neuroinflammation* 2005; 2: 23.
- [23] Billings L, Oddo S, Green K, McGaugh J and Laferla F. Intraneuronal Abeta causes the onset of early Alzheimer's disease-related cognitive deficits in transgenic mice. *Neuron* 2005; 45: 675 - 688.
- [24] Junker A, Krumbholz M, Eisele S, Mohan H, Augstein F, Bittner R, Lassmann H, Wekerle H, Hohlfeld R and Meinel E. MicroRNA profiling of multiple sclerosis lesions identifies modulators of the regulatory protein CD47. *Brain* 2009; 132: 3342-3352.
- [25] FARAONI, #160, Isabella, ANTONETTI R, Francesca, CARDONE, John, BONMASSAR and Enzo. miR-155 gene: A typical multifunctional microRNA. Amsterdam, PAYS-BAS: Elsevier, 2009.
- [26] Buck AH, Perot J, Chisholm MA, Kumar DS, Tuddenham L, Cognat V, Marcinowski L, Dölken L and Pfeffer S. Post-transcriptional regulation of miR-27 in murine cytomegalovirus infection. *RNA* 2010; 16: 307-315.
- [27] Johnson CD, Esquela-Kerscher A, Stefani G, Byrom M, Kelnar K, Ovcharenko D, Wilson M, Wang X, Shelton J, Shingara J, Chin L, Brown D and Slack FJ. The let-7 MicroRNA Represses Cell Proliferation Pathways in Human Cells. *Cancer Research* 2007; 67: 7713-7722.
- [28] Schultz J, Lorenz, Peter, Gross, Gerd, Ibrahim, Saleh and Kunz, Manfred. MicroRNA let-7b targets important cell cycle molecules in malignant melanoma cells and interferes with anchorage-independent growth. *Cell Research* 2008; 18: 549-557.
- [29] Chhabra R, Adlakha YK, Hariharan M, Scaria V and Saini N. Upregulation of miR-23a~27a~24-2 Cluster Induces Caspase-Dependent and -Independent Apoptosis in Human Embryonic Kidney Cells. *PLoS ONE* 2009; 4: e5848.
- [30] Morris JC, Edland S, Clark C, Galasko D, Koss E, Mohs R, van Belle G, Fillenbaum G and Heyman A. The Consortium to Establish a Registry for Alzheimer's Disease (CERAD): Part IV. Rates of cognitive change in the longitudinal assessment of probable Alzheimer's disease. *Neurology* 1993; 43: 2457-.
- [31] Kenchappa P, Yadav A, Singh G, Nandana S and Banerjee K. Rescue of TNF α -inhibited neuronal cells by IGF-1 involves Akt and c-Jun N-terminal kinases. *Journal of Neuroscience Research* 2004; 76: 466-474.
- [32] Livak KJ and Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2⁻(-Delta Delta C(T)) Method. *Methods (Duluth)* 2001; 25: 402-408.
- [33] Beutler B and Cerami A. The Biology of Cachectin/TNF -- A Primary Mediator of the Host Response. *Annual Review of Immunology* 1989; 7: 625-655.
- [34] Deleault KM, Skinner SJ and Brooks SA. Tristetraprolin regulates TNF TNF-[alpha] mRNA stability via a proteasome dependent mechanism involving the combined action of the ERK and p38 pathways. *Molecular Immunology* 2008; 45: 13-24.
- [35] Lanzrein A, Johnston C, Perry V, Jobst K, King E and Smith A. Longitudinal study of inflammatory factors in serum, cerebrospinal fluid, and brain tissue in Alzheimer disease: interleukin-1beta, interleukin-6, interleukin-1 receptor antagonist, tumor necrosis factor-alpha, the soluble tumor necrosis factor receptors I and II, and alpha1-antichymotrypsin. *Alzheimer Dis Assoc Disord* 1998; 12: 215 - 227.
- [36] Lukiw WJ. Micro-RNA speciation in fetal, adult and Alzheimer's disease hippocampus. *NeuroReport* 2007; 18: 297-300 210.1097/WNR.1090b1013e3280148e3280148b.
- [37] Schonrock N, Ke YD, Humphreys D, Staufienbiel M, Ittner LM, Preiss T and Götz J. Neuronal MicroRNA Deregulation in Response to Alzheimer's Disease Amyloid- β . *PLoS ONE* 2010; 5: e11070.
- [38] Smirnova L, Gräfe A, Seiler A, Schumacher S, Nitsch R and Wulczyn FG. Regulation of miRNA expression during neural cell specification. *European Journal of Neuroscience* 2005; 21:

- 1469-1477.
- [39] Carrettiero DC, Hernandez I, Neveu P, Pagiannakopoulos T and Kosik KS. The Cochaperone BAG2 Sweeps Paired Helical Filament-Insoluble Tau from the Microtubule. *J. Neurosci.* 2009; 29: 2151-2161.
- [40] Kolachala VL, Wang L, Obertone TS, Prasad M, Yan Y, Dalmaso G, Gewirtz AT, Merlin D and Sitaraman SV. Adenosine 2B Receptor Expression Is Post-transcriptionally Regulated by MicroRNA. *Journal of Biological Chemistry* 2010; 285: 18184-18190.
- [41] Cullen BR. Transcription and Processing of Human microRNA Precursors. *Molecular Cell* 2004; 16: 861-865.
- [42] Evangelisti C, Florian MC, Massimi I, Dominici C, Giannini G, Galardi S, Buè MC, Massalini S, McDowell HP, Messi E, Gulino A, Farace MG and Ciafrè SA. MiR-128 up-regulation inhibits Reelin and DCX expression and reduces neuroblastoma cell motility and invasiveness. *The FASEB Journal* 2009; 23: 4276-4287.
- [43] Fukuda Y, Kawasaki H and Taira K. Exploration of human miRNA target genes in neuronal differentiation. *Nucleic Acids Symposium Series* 2005; 49: 341-342.
- [44] Tili E, Michaille J-J, Cimino A, Costinean S, Dumitru CD, Adair B, Fabbri M, Alder H, Liu CG, Calin GA and Croce CM. Modulation of miR-155 and miR-125b Levels following Lipopolysaccharide/TNF- α Stimulation and Their Possible Roles in Regulating the Response to Endotoxin Shock. *The Journal of Immunology* 2007; 179: 5082-5089.
- [45] O'Connell RM, Rao DS, Chaudhuri AA, Boldin MP, Taganov KD, Nicoll J, Paquette RL and Baltimore D. Sustained expression of microRNA -155 in hematopoietic stem cells causes a myeloproliferative disorder. *The Journal of Experimental Medicine* 2008; 205: 585-594.
- [46] Kehoe PG, Miners S and Love S. Angiotensins in Alzheimer's disease friend or foe? *Trends in neurosciences* 2009; 32: 619-628.
- [47] Harold D, Abraham R, Hollingworth P, Sims R, Gerrish A, Hamshere ML, Pahwa JS, Moskvina V, Dowzell K, Williams A, Jones N, Thomas C, Stretton A, Morgan AR, Lovestone S, Powell J, Proitsi P, Lupton MK, Brayne C, Rubinsztein DC, Gill M, Lawlor B, Lynch A, Morgan K, Brown KS, Passmore PA, Craig D, McGuinness B, Todd S, Holmes C, Mann D, Smith AD, Love S, Kehoe PG, Hardy J, Mead S, Fox N, Rossor M, Collinge J, Maier W, Jessen F, Schurmann B, van den Bussche H, Heuser I, Kornhuber J, Wiltfang J, Dichgans M, Frolich L, Hampel H, Hull M, Rujescu D, Goate AM, Kauwe JSK, Cruchaga C, Nowotny P, Morris JC, Mayo K, Sleegers K, Bettens K, Engelborghs S, De Deyn PP, Van Broeckhoven C, Livingston G, Bass NJ, Gurling H, McQuillin A, Gwilliam R, Deloukas P, Al-Chalabi A, Shaw CE, Tsolaki M, Singleton AB, Guerreiro R, Muhleisen TW, Nothen MM, Moebus S, Jockel K-H, Klopp N, Wichmann HE, Carrasquillo MM, Pankratz VS, Younkin SG, Holmans PA, O'Donovan M, Owen MJ and Williams J. Genome-wide association study identifies variants at CLU and PICALM associated with Alzheimer's disease. *Nat Genet* 2009; 41: 1088-1093.
- [48] Vasudevan S, Tong Y and Steitz JA. Switching from Repression to Activation: MicroRNAs Can Up-Regulate Translation. *Science* 2007; 318: 1931-1934.